

References

- Acacio GL, Barini R, Pinto Junior W, *et al.* 2001. Nuchal translucency: an ultrasound marker for fetal chromosomal abnormalities. *Sao Paulo Med J*, **119**: 19-23.
- Achiron R, Barkai G, Katznelson MB, *et al.* 1991. Fetal lateral ventricle choroid plexus cysts: the dilemma of amniocentesis. *Obstet Gynecol*, **78**: 815-818.
- ACOG. 2007a. ACOG Practice Bulletin No. 77: screening for fetal chromosomal abnormalities. *Obstet Gynecol*, **109**: 217-227.
- ACOG. 2007b. ACOG Practice Bulletin No. 88, December 2007. Invasive prenatal testing for aneuploidy. *Obstet Gynecol*, **110**: 1459-1467.
- Aitken DA, McCaw G, Crossley JA, *et al.* 1993. First-trimester biochemical screening for fetal chromosome abnormalities and neural tube defects. *Prenat Diagn*, **13**: 681-689.
- Aitken DA, Wallace EM, Crossley JA, *et al.* 1996. Dimeric inhibin A as a marker for Down's syndrome in early pregnancy. *N Engl J Med*, **334**: 1231-1236.
- Akolekar R, Perez Penco JM, Skyfta E, *et al.* 2010. Maternal serum placental protein 13 at eleven to thirteen weeks in chromosomally abnormal pregnancies. *Fetal Diagn Ther*, **27**: 72-77.
- Alfirevic Z, Sundberg K, Brigham S. 2008. Amniocentesis and chorionic villus sampling for prenatal diagnosis. *Cochrane Database Syst Rev*. CD003252.
- Ammala P, Hiilesmaa VK, Liukkonen S, *et al.* 1993. Randomized trial comparing first-trimester transcervical chorionic villus sampling and second-trimester amniocentesis. *Prenat Diagn*, **13**: 919-927.
- Anandakumar C, Chew S, Wong YC, *et al.* 1999. The sensitivity of the trivariate analysis using maternal serum alpha-feto protein, human chorionic gonadotrophin and maternal age in screening for fetal aneuploidy in mothers above the age of 35. *J Perinat Med*, **27**: 178-182.
- Antolin E, Comas C, Torrents M, *et al.* 2001. The role of ductus venosus blood flow assessment in screening for chromosomal abnormalities at 10-16 weeks of gestation. *Ultrasound Obstet Gynecol*, **17**: 295-300.
- Arab H, Siegel-Bartelt J, Wong PY, Doran T. 1988. Maternal serum beta human chorionic gonadotropin (MSHCG) combined with maternal serum alpha-fetoprotein (MSAFP) appears superior for prenatal screening for Down syndrome (DS) than entire test alone. *Am J Hum Genet*, **43**: a225.
- Ashwood ER, Cheng E, Luthy DA. 1987. Maternal serum alpha-fetoprotein and fetal trisomy-21 in women 35 years and older: implications for alpha-fetoprotein screening programs. *Am J Med Genet*, **26**: 531-539.

- Bahado-Singh RO, Choi SJ, Oz U, *et al.* 2003. Early second-trimester individualized estimation of trisomy 18 risk by ultrasound. *Obstet Gynecol*, **101**: 463-468.
- Bahado-Singh RO, Lynch L, Deren O, *et al.* 1997. First-trimester growth restriction and fetal aneuploidy: the effect of type of aneuploidy and gestational age. *Am J Obstet Gynecol*, **176**: 976-980.
- Bakos O, Moen KS, Hansson S. 1998. Prenatal karyotyping of choroid plexus cysts. *Eur J Ultrasound*, **8**: 79-83.
- Barkai G, Goldman B, Ries L, *et al.* 1993. Expanding multiple marker screening for Down's syndrome to include Edward's syndrome. *Prenat Diagn*, **13**: 843-850.
- Bartels I, Thiele M, Bogart MH. 1990. Maternal serum hCG and SP1 in pregnancies with fetal aneuploidy. *Am J Med Genet*, **37**: 261-264.
- Baty BJ, Blackburn BL, Carey JC. 1994a. Natural history of trisomy 18 and trisomy 13: I. Growth, physical assessment, medical histories, survival, and recurrence risk. *Am J Med Genet*, **49**: 175-188.
- Baty BJ, Jorde LB, Blackburn BL, *et al.* 1994b. Natural history of trisomy 18 and trisomy 13: II. Psychomotor development. *Am J Med Genet*, **49**: 189-194.
- Beke A, Barakonyi E, Belics Z, *et al.* 2008. Risk of chromosome abnormalities in the presence of bilateral or unilateral choroid plexus cysts. *Fetal Diagn Ther*, **23**: 185-191.
- Bekker MN, Twisk JW, van Vugt JM. 2004. Reproducibility of the fetal nasal bone length measurement. *J Ultrasound Med*, **23**: 1613-1618.
- Benacerraf BR, Barss VA, Laboda LA. 1985. A sonographic sign for the detection in the second trimester of the fetus with Down's syndrome. *Am J Obstet Gynecol*, **151**: 1078-1079.
- Benacerraf BR, Cnann A, Gelman R, *et al.* 1989. Can sonographers reliably identify anatomic features associated with Down syndrome in fetuses? *Radiology*, **173**: 377-380.
- Benacerraf BR, Harlow B, Frigoletto FD, Jr. 1990. Are choroid plexus cysts an indication for second-trimester amniocentesis? *Am J Obstet Gynecol*, **162**: 1001-1006.
- Benacerraf BR, Miller WA, Frigoletto FD, Jr. 1988. Sonographic detection of fetuses with trisomies 13 and 18: accuracy and limitations. *Am J Obstet Gynecol*, **158**: 404-409.
- Benacerraf BR, Nadel A, Bromley B. 1994. Identification of second-trimester fetuses with autosomal trisomy by use of a sonographic scoring index. *Radiology*, **193**: 135-140.
- Benacerraf BR, Neuberg D, Bromley B, *et al.* 1992. Sonographic scoring index for prenatal detection of chromosomal abnormalities. *J Ultrasound Med*, **11**: 449-458.
- Benn PA. 2010. Prenatal Diagnosis of Chromosomal Abnormalities through Amniocentesis. In *Genetic Disorders and the Fetus: Diagnosis, Prevention and Treatment*, Milunsky A, Milunsky JM (eds.). 6th ed. Wiley-Blackwell: West Sussex; 194-272.

- Benn PA, Horne D, Craffey A, *et al.* 1996. Maternal serum screening for birth defects: results of a Connecticut regional program. *Conn Med*, **60**: 323-327.
- Benn PA, Leo MV, Rodis JF, *et al.* 1999. Maternal serum screening for fetal trisomy 18: a comparison of fixed cutoff and patient-specific risk protocols. *Obstet Gynecol*, **93**: 707-711.
- Bersinger NA, Brizot ML, Johnson A, *et al.* 1994. First trimester maternal serum pregnancy-associated plasma protein A and pregnancy-specific beta 1-glycoprotein in fetal trisomies. *Br J Obstet Gynaecol*, **101**: 970-974.
- Bersinger NA, Leporrier N, Herrou M, *et al.* 1999. Maternal serum pregnancy-associated plasma protein A (PAPP-A) but not pregnancy-specific beta1-glycoprotein (SP1) is a useful second-trimester marker for fetal trisomy 18. *Prenat Diagn*, **19**: 537-541.
- Bersu ET, Ramirez-Castro JL. 1977. Anatomical analysis of the developmental effects of aneuploidy in man--the 18-trisomy syndrome: I. Anomalies of the head and neck. *Am J Med Genet*, **1**: 173-193.
- Bestwick JP, Huttly WJ, Wald NJ. 2010. Distribution of nuchal translucency in antenatal screening for Down's syndrome. *J Med Screen*, **17**: 8-12.
- Bewley S, Roberts LJ, Mackinson AM, *et al.* 1995. First trimester fetal nuchal translucency: problems with screening the general population. 2. *Br J Obstet Gynaecol*, **102**: 386-388.
- Biagiotti R, Cariati E, Brizzi L, *et al.* 1998. Maternal serum screening for trisomy 18 in the first trimester of pregnancy. *Prenat Diagn*, **18**: 907-913.
- Bianchi DW, Simpson JL, Jackson LG, *et al.* 2002. Fetal gender and aneuploidy detection using fetal cells in maternal blood: analysis of NIFTY I data. National Institute of Child Health and Development Fetal Cell Isolation Study. *Prenat Diagn*, **22**: 609-615.
- Bilardo CM, Muller MA, Zikulnig L, *et al.* 2001. Ductus venosus studies in fetuses at high risk for chromosomal or heart abnormalities: relationship with nuchal translucency measurement and fetal outcome. *Ultrasound Obstet Gynecol*, **17**: 288-294.
- Bindra R, Heath V, Liao A, *et al.* 2002. One-stop clinic for assessment of risk for trisomy 21 at 11-14 weeks: a prospective study of 15 030 pregnancies. *Ultrasound Obstet Gynecol*, **20**: 219-225.
- Bird LM, Dixon B, Masser-Frye D, *et al.* 2002. Choroid plexus cysts in the mid-trimester fetus--practical application suggests superiority of an individualized risk method of counseling for trisomy 18. *Prenat Diagn*, **22**: 792-797.
- Blitzer MG, Carmi R, Blakemore K, Andrews J, Romen I, Schwartz S. 1991. Low maternal serum human chorionic gonadotropin (MS-hCG) in second trimester trisomy 18 pregnancies. *Am J Hum Genet*, **49**: a211.
- Bogart MH, Golbus MS, Sorg ND, *et al.* 1989. Human chorionic gonadotropin levels in pregnancies with aneuploid fetuses. *Prenat Diagn*, **9**: 379-384.

- Bogart MH, Pandian MR, Jones OW. 1987. Abnormal maternal serum chorionic gonadotropin levels in pregnancies with fetal chromosome abnormalities. *Prenat Diagn*, **7**: 623-630.
- Borrell A, Casals E, Fortuny A, *et al.* 2004. First-trimester screening for trisomy 21 combining biochemistry and ultrasound at individually optimal gestational ages. An interventional study. *Prenat Diagn*, **24**: 541-545.
- Borrell A, Costa D, Martinez JM, *et al.* 1997. Criteria for fetal nuchal thickness cut-off: a re-evaluation. *Prenat Diagn*, **17**: 23-29.
- Borrell A, Martinez JM, Seres A, *et al.* 2003. Ductus venosus assessment at the time of nuchal translucency measurement in the detection of fetal aneuploidy. *Prenat Diagn*, **23**: 921-926.
- Borrell A, Perez M, Figueras F, *et al.* 2007. Reliability analysis on ductus venosus assessment at 11-14 weeks' gestation in a high-risk population. *Prenat Diagn*, **27**: 442-446.
- Bottalico JN, Chen X, Tartaglia M, *et al.* 2009. Second-trimester genetic sonogram for detection of fetal chromosomal abnormalities in a community-based antenatal testing unit. *Ultrasound Obstet Gynecol*, **33**: 161-168.
- Bradley LA, Horowitz JA, Dowman AC, Ponting NR, Peterson LM. 1994. Triple marker screening for fetal Down syndrome. *International Journal of Pediatrics*, **9**: 168-174.
- Brambati B, Lanzani A, Tului L. 1990. Transabdominal and transcervical chorionic villus sampling: efficiency and risk evaluation of 2,411 cases. *Am J Med Genet*, **35**: 160-164.
- Brambati B, Tului L, Bonacchi I, *et al.* 1994. Serum PAPP-A and free beta-hCG are first-trimester screening markers for Down syndrome. *Prenat Diagn*, **14**: 1043-1047.
- Breathnach FM, Malone FD, Lambert-Messerlian G, *et al.* 2007. First- and second-trimester screening: detection of aneuploidies other than Down syndrome. *Obstet Gynecol*, **110**: 651-657.
- Brewer CM, Holloway SH, Stone DH, Carothers AD, FitzPatrick DR. 2002. Survival in trisomy 13 and trisomy 18 cases ascertained from population based registers. *J Med Genet*, **39**: e54.
- Brizot ML, Carvalho MH, Liao AW, *et al.* 2001. First-trimester screening for chromosomal abnormalities by fetal nuchal translucency in a Brazilian population. *Ultrasound Obstet Gynecol*, **18**: 652-655.
- Brizot ML, Snijders RJ, Bersinger NA, *et al.* 1994. Maternal serum pregnancy-associated plasma protein A and fetal nuchal translucency thickness for the prediction of fetal trisomies in early pregnancy. *Obstet Gynecol*, **84**: 918-922.
- Brizot ML, Snijders RJ, Butler J, *et al.* 1995. Maternal serum hCG and fetal nuchal translucency thickness for the prediction of fetal trisomies in the first trimester of pregnancy. *Br J Obstet Gynaecol*, **102**: 127-132.

- Brock DJ, Bolton AE, Monaghan JM. 1973. Prenatal diagnosis of anencephaly through maternal serum-alpha-fetoprotein measurement. *Lancet*, **2**: 923-924.
- Brock DJ, Sutcliffe RG. 1972. Alpha-fetoprotein in the antenatal diagnosis of anencephaly and spina bifida. *Lancet*, **2**: 197-199.
- Bromley B, Doubilet P, Frigoletto FD, Jr., et al. 1994. Is fetal hyperechoic bowel on second-trimester sonogram an indication for amniocentesis? *Obstet Gynecol*, **83**: 647-651.
- Bronshtein M, Rottem S, Yoffe N, et al. 1989. First-trimester and early second-trimester diagnosis of nuchal cystic hygroma by transvaginal sonography: diverse prognosis of the septated from the nonseptated lesion. *Am J Obstet Gynecol*, **161**: 78-82.
- Bronsteen R, Lee W, Vettraino IM, et al. 2004. Second-trimester sonography and trisomy 18. *J Ultrasound Med*, **23**: 233-240.
- Brumfield CG, Wenstrom KD, Owen J, et al. 2000. Ultrasound findings and multiple marker screening in trisomy 18. *Obstet Gynecol*, **95**: 51-54.
- Bundy AL, Saltzman DH, Pober B, et al. 1986. Antenatal sonographic findings in trisomy 18. *J Ultrasound Med*, **5**: 361-364.
- Burton BK, Prins GS, Verp MS. 1993. A prospective trial of prenatal screening for Down syndrome by means of maternal serum alpha-fetoprotein, human chorionic gonadotropin, and unconjugated estriol. *Am J Obstet Gynecol*, **169**: 526-530.
- Burton BK, Schulz CJ, Burd LI. 1992. Limb anomalies associated with chorionic villus sampling. *Obstet Gynecol*, **79**: 726-730.
- Burton BK, Sowers SG, Nelson LH. 1983. Maternal serum alpha-fetoprotein screening in North Carolina: experience with more than twelve thousand pregnancies. *Am J Obstet Gynecol*, **146**: 439-444.
- Byrne J, Warburton D, Kline J, et al. 1985. Morphology of early fetal deaths and their chromosomal characteristics. *Teratology*, **32**: 297-315.
- Camurri L, Ventura A. 1989. Prospective study on trisomy 18 and fetal choroid plexus cysts. *Prenat Diagn*, **9**: 742.
- Canick JA, Palomaki GE, Osathanondh R. 1990. Prenatal screening for trisomy 18 in the second trimester. *Prenat Diagn*, **10**: 546-548.
- Carey JC. 2005. Trisomy 18 and trisomy 13 syndrome. In *In: Management of Genetic Syndromes*, (eds.). 2nd edition ed. Wiley-Liss: New York; 555-568.
- Carey JC. 2006. Introductory comments special section: Trisomy 18. *Am J Med Genet Part A*, **140A**: 935-936.
- Carter PE, Pearn JH, Bell J, et al. 1985. Survival in trisomy 18. Life tables for use in genetic counselling and clinical paediatrics. *Clin Genet*, **27**: 59-61.

- CEMAT. 1998. Randomised trial to assess safety and fetal outcome of early and midtrimester amniocentesis. The Canadian Early and Mid-trimester Amniocentesis Trial (CEMAT) Group. *Lancet*, **351**: 242-247.
- Chan L, Hixson JL, Laifer SA, *et al.* 1989. A sonographic and karyotypic study of second-trimester fetal choroid plexus cysts. *Obstet Gynecol*, **73**: 703-706.
- Chao AS, Chung CL, Wu CD, *et al.* 1999. Second trimester maternal serum screening using alpha fetoprotein, free beta human chorionic gonadotropin and maternal age specific risk: result of chromosomal abnormalities detected in screen positive for Down syndrome in an Asian population. *Acta Obstet Gynecol Scand*, **78**: 393-397.
- Chasen ST, Sharma G, Kalish RB, *et al.* 2003. First-trimester screening for aneuploidy with fetal nuchal translucency in a United States population. *Ultrasound Obstet Gynecol*, **22**: 149-151.
- Chen XQ, Stroun M, Magnenat JL, *et al.* 1996. Microsatellite alterations in plasma DNA of small cell lung cancer patients. *Nat Med*, **2**: 1033-1035.
- Cheng PJ, Liu CM, Chueh HY, *et al.* 2003. First-trimester nuchal translucency measurement and echocardiography at 16 to 18 weeks of gestation in prenatal detection for trisomy 18. *Prenat Diagn*, **23**: 248-251.
- Cheng PJ, Shaw SW, Soong YK. 2006. Association of fetal choroid plexus cysts with trisomy 18 in a population previously screened by nuchal translucency thickness measurement. *J Soc Gynecol Investig*, **13**: 280-284.
- Chinn DH, Miller EI, Worthy LM, *et al.* 1991. Sonographically detected fetal choroid plexus cysts. Frequency and association with aneuploidy. *J Ultrasound Med*, **10**: 255-258.
- Chitkara U, Cogswell C, Norton K, *et al.* 1988. Choroid plexus cysts in the fetus: a benign anatomic variant or pathologic entity? Report of 41 cases and review of the literature. *Obstet Gynecol*, **72**: 185-189.
- Chitty LS, Chudleigh P, Wright E, *et al.* 1998. The significance of choroid plexus cysts in an unselected population: results of a multicenter study. *Ultrasound Obstet Gynecol*, **12**: 391-397.
- Chiu RW, Chan KC, Gao Y, *et al.* 2008. Noninvasive prenatal diagnosis of fetal chromosomal aneuploidy by massively parallel genomic sequencing of DNA in maternal plasma. *Proc Natl Acad Sci U S A*, **105**: 20458-20463.
- Cho RC, Chu P, Smith-Bindman R. 2009. Second trimester prenatal ultrasound for the detection of pregnancies at increased risk of Trisomy 18 based on serum screening. *Prenat Diagn*, **29**: 129-139.
- Chudleigh P, Pearce JM, Campbell S. 1984. The prenatal diagnosis of transient cysts of the fetal choroid plexus. *Prenat Diagn*, **4**: 135-137.

- Cicero S, Avgidou K, Rembouskos G, *et al.* 2006. Nasal bone in first-trimester screening for trisomy 21. *Am J Obstet Gynecol*, **195**: 109-114.
- Cicero S, Curcio P, Papageorghiou A, *et al.* 2001. Absence of nasal bone in fetuses with trisomy 21 at 11-14 weeks of gestation: an observational study. *Lancet*, **358**: 1665-1667.
- Cicero S, Longo D, Rembouskos G, *et al.* 2003. Absent nasal bone at 11-14 weeks of gestation and chromosomal defects. *Ultrasound Obstet Gynecol*, **22**: 31-35.
- Cicero S, Rembouskos G, Vandecruys H, *et al.* 2004. Likelihood ratio for trisomy 21 in fetuses with absent nasal bone at the 11-14-week scan. *Ultrasound Obstet Gynecol*, **23**: 218-223.
- Cicero S, Spencer K, Avgidou K, *et al.* 2005. Maternal serum biochemistry at 11-13(+6) weeks in relation to the presence or absence of the fetal nasal bone on ultrasonography in chromosomally abnormal fetuses: an updated analysis of integrated ultrasound and biochemical screening. *Prenat Diagn*, **25**: 977-983.
- Clark SL, DeVore GR, Sabey PL. 1988. Prenatal diagnosis of cysts of the fetal choroid plexus. *Obstet Gynecol*, **72**: 585-587.
- Coco C, Jeanty P. 2004. Karyotyping of fetuses with isolated choroid plexus cysts is not justified in an unselected population. *J Ultrasound Med*, **23**: 899-906.
- Costa JM, Benachi A, Gautier E, *et al.* 2001. First-trimester fetal sex determination in maternal serum using real-time PCR. *Prenat Diagn*, **21**: 1070-1074.
- Cowchock FS, Rush DA. 1984. Low maternal serum AFP and Down syndrome. *Lancet*, **ii**: 161.
- Crider KS, Olney RS, Cragan JD. 2008. Trisomies 13 and 18: population prevalences, characteristics, and prenatal diagnosis, metropolitan Atlanta, 1994-2003. *Am J Med Genet A*, **146**: 820-826.
- Crossley JA, Aitken DA, Connor JM. 1991. Prenatal screening for chromosome abnormalities using maternal serum chorionic gonadotrophin, alpha-fetoprotein, and age. *Prenat Diagn*, **11**: 83-101.
- Crossley JA, Aitken DA, Connor JM. 1993. Second-trimester unconjugated oestriol levels in maternal serum from chromosomally abnormal pregnancies using an optimized assay. *Prenat Diagn*, **13**: 271-280.
- Cuckle H. 2010. Monitoring quality control of nuchal translucency. *Clin Lab Med*, **30**: 593-604.
- Cuckle HS, Wald NJ, Lindenbaum RH. 1984. Maternal serum alpha-fetoprotein measurement: a screening test for Down syndrome. *Lancet*, **1**: 926-929.
- Cuckle HS, Wald NJ, Thompson SG. 1987. Estimating a woman's risk of having a pregnancy associated with Down's syndrome using her age and serum alpha-fetoprotein level. *Br J Obstet Gynaecol*, **94**: 387-402.

- Cullen MT, Gabrielli S, Green JJ, *et al.* 1990. Diagnosis and significance of cystic hygroma in the first trimester. *Prenat Diagn*, **10**: 643-651.
- Cullen MT, Green JJ, Scioscia AL, *et al.* 1995. Ultrasonography in the detection of aneuploidy in the first trimester. *J Ultrasound Med*, **14**: 559-563.
- Darnule AT, Greenberg F, Warrington RE, *et al.* 1990. Superoxide dismutase levels in amniotic fluid. *Prenat Diagn*, **10**: 757-758.
- Darnule AT, Schmidt D, Weyland B, Greenberg E, Rose E, Alpert E. 1990. Serum hCG, AFP and unconjugated estriol levels in trisomy 18 in mid-trimester. *Am J Hum Genet*, **47**: a272.
- De Souza E, Halliday J, Chan A, *et al.* 2009. Recurrence risks for trisomies 13, 18, and 21. *Am J Med Genet A*, **149A**: 2716-2722.
- DeRoo TR, Harris RD, Sargent SK, *et al.* 1988. Fetal choroid plexus cysts: prevalence, clinical significance, and sonographic appearance. *AJR Am J Roentgenol*, **151**: 1179-1181.
- DeVore GR. 2000. Second trimester ultrasonography may identify 77 to 97% of fetuses with trisomy 18. *J Ultrasound Med*, **19**: 565-576.
- Dhaifalah I, Santavy J, Zapletalova J. 2006. Screening for chromosomal anomalies in the first trimester: a report on the first year of prospective screening for chromosomal anomalies in the first trimester in the Czech Republic. *Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub*, **150**: 275-278.
- Dicke JM, Gray DL, Songster GS, *et al.* 1989. Fetal biometry as a screening tool for the detection of chromosomally abnormal pregnancies. *Obstet Gynecol*, **74**: 726-729.
- Digiovanni LM, Quinlan MP, Verp MS. 1997. Choroid plexus cysts: infant and early childhood developmental outcome. *Obstet Gynecol*, **90**: 191-194.
- DiMaio MS, Baumgarten A, Greenstein RM, *et al.* 1987. Screening for fetal Down's syndrome in pregnancy by measuring maternal serum alpha-fetoprotein levels. *N Engl J Med*, **317**: 342-346.
- Dix U, Gams M, Grubisic A, *et al.* 1988. [Alpha fetoprotein in maternal serum in pregnancies with trisomy 18]. *Z Geburtshilfe Perinatol*, **192**: 231-233.
- Doran TA, Cadesky K, Wong PY, *et al.* 1986. Maternal serum alpha-fetoprotein and fetal autosomal trisomies. *Am J Obstet Gynecol*, **154**: 277-281.
- Down JLH. 1866. Observations on an ethnic classification of idiots. *London Hospital Reports*, **3**: 259-262.
- Droste S, FitzSimmons J, Pascoe-Mason J, *et al.* 1990. Growth of linear parameters in trisomy 18 fetuses. *Am J Obstet Gynecol*, **163**: 158-161.
- Droste S, Hendricks SK, Von Alfrey H, *et al.* 1991. Cystic hygroma colli: perinatal outcome after prenatal diagnosis. *J Perinat Med*, **19**: 449-454.

- Drugan A, Johnson MP, Reichler A, *et al.* 1996. Second-trimester minor ultrasound anomalies: impact on the risk of aneuploidy associated with advanced maternal age. *Obstet Gynecol*, **88**: 203-206.
- Duval S, Tweedie R. 2000. Trim and fill: A simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics*, **56**: 455-463.
- Edwards JH, Harnden DG, Cameron AH, *et al.* 1960. A new trisomic syndrome. *Lancet*, **1**: 787-790.
- Embleton ND, Wyllie JP, Wright MJ, *et al.* 1996. Natural history of trisomy 18. *Arch Dis Child Fetal Neonatal Ed*, **75**: F38-41.
- Faiola S, Tsoi E, Huggon IC, *et al.* 2005. Likelihood ratio for trisomy 21 in fetuses with tricuspid regurgitation at the 11 to 13 + 6-week scan. *Ultrasound Obstet Gynecol*, **26**: 22-27.
- Falcon O, Faiola S, Huggon I, *et al.* 2006. Fetal tricuspid regurgitation at the 11 + 0 to 13 + 6-week scan: association with chromosomal defects and reproducibility of the method. *Ultrasound Obstet Gynecol*, **27**: 609-612.
- Falcon O, Wegrzyn P, Faro C, *et al.* 2005. Gestational sac volume measured by three-dimensional ultrasound at 11 to 13 + 6 weeks of gestation: relation to chromosomal defects. *Ultrasound Obstet Gynecol*, **25**: 546-550.
- Fan HC, Blumenfeld YJ, Chitkara U, *et al.* 2008. Noninvasive diagnosis of fetal aneuploidy by shotgun sequencing DNA from maternal blood. *Proc Natl Acad Sci U S A*, **105**: 16266-16271.
- Ferguson-Smith MA, Rawlinson HA, May HM, *et al.* 1978. Avoidance of anencephalic and spina bifida births by maternal serum-alpha-fetoprotein screening. *Lancet*, **1**: 1330-1333.
- Feuchtbaum LB, Currier RJ, Lorey FW, *et al.* 2000. Prenatal ultrasound findings in affected and unaffected pregnancies that are screen-positive for trisomy 18: the California experience. *Prenat Diagn*, **20**: 293-299.
- Field B, Mitchell G, Kerr C. 1976. Antenatal diagnosis of neural tube defects. *Med J Aust*, **2**: 5-7.
- Firth HV, Boyd PA, Chamberlain P, *et al.* 1991. Limb abnormalities and chorion villus sampling. *Lancet*, **338**: 51.
- FitzPatrick DR, Boyd E. 1989. Recurrences of trisomy 18 and trisomy 13 after trisomy 21. *Hum Genet*, **82**: 301.
- Fitzsimmons J, Wilson D, Pascoe-Mason J, *et al.* 1989. Choroid plexus cysts in fetuses with trisomy 18. *Obstet Gynecol*, **73**: 257-260.
- Forrester MB, Merz RD. 1999. Trisomies 13 and 18: prenatal diagnosis and epidemiologic studies in Hawaii, 1986-1997. *Genet Test*, **3**: 335-340.

- Forrester MB, Merz RD. 2002. Pregnancy outcome distribution and prenatal diagnosis of autosomal abnormalities, Hawaii, 1986-1999. *Teratology*, **66 Suppl 1**: S7-11.
- Gabrielli S, Reece EA, Pilu G, et al. 1989. The clinical significance of prenatally diagnosed choroid plexus cysts. *Am J Obstet Gynecol*, **160**: 1207-1210.
- Gardener RJM, Sutherland GR. 1996. Chromosome abnormalities and genetic counseling. Oxford Monograph on Medical Genetics. No. 29. 2nd Edition. Oxford: Oxford University Press.
- Geary M, Patel S, Lamont R. 1997. Isolated choroid plexus cysts and association with fetal aneuploidy in an unselected population. *Ultrasound Obstet Gynecol*, **10**: 171-173.
- Gembruch U, Hansmann M, Bald R, et al. 1988. Prenatal diagnosis and management in fetuses with cystic hygromata colli. *Eur J Obstet Gynecol Reprod Biol*, **29**: 241-255.
- Ghidini A, Strobelt N, Locatelli A, et al. 2000. Isolated fetal choroid plexus cysts: role of ultrasonography in establishment of the risk of trisomy 18. *Am J Obstet Gynecol*, **182**: 972-977.
- Ginsberg N, Cadkin A, Pergament E, et al. 1990. Ultrasonographic detection of the second-trimester fetus with trisomy 18 and trisomy 21. *Am J Obstet Gynecol*, **163**: 1186-1190.
- Goetzinger KR, Stamilio DM, Dicke JM, et al. 2008. Evaluating the incidence and likelihood ratios for chromosomal abnormalities in fetuses with common central nervous system malformations. *Am J Obstet Gynecol*, **199**: 285 e281-286.
- Golden CM, Ryan LM, Holmes LB. 2003. Chorionic villus sampling: a distinctive teratogenic effect on fingers? *Birth Defects Res A Clin Mol Teratol*, **67**: 557-562.
- Goldstein H, Nielsen KG. 1988. Rates and survival of individuals with trisomy 13 and 18. Data from a 10-year period in Denmark. *Clin Genet*, **34**: 366-372.
- Gonzalez R, Silva JM, Sanchez A, et al. 2000. Microsatellite alterations and TP53 mutations in plasma DNA of small-cell lung cancer patients: follow-up study and prognostic significance. *Ann Oncol*, **11**: 1097-1104.
- Grandjean H, Larroque D, Levi S. 1998. Detection of chromosomal abnormalities, an outcome of ultrasound screening. The Eurofetus Team. *Ann N Y Acad Sci*, **847**: 136-140.
- Gray DL, Winborn RC, Suessen TL, et al. 1996. Is genetic amniocentesis warranted when isolated choroid plexus cysts are found? *Prenat Diagn*, **16**: 983-990.
- Greenberg F, Schmidt D, Darnule AT, et al. 1992. Maternal serum alpha-fetoprotein, beta-human chorionic gonadotropin, and unconjugated estriol levels in midtrimester trisomy 18 pregnancies. *Am J Obstet Gynecol*, **166**: 1388-1392.
- Gross SJ, Shulman LP, Tolley EA, et al. 1995. Isolated fetal choroid plexus cysts and trisomy 18: a review and meta-analysis. *Am J Obstet Gynecol*, **172**: 83-87.

- Gupta JK, Cave M, Lilford RJ, *et al.* 1995. Clinical significance of fetal choroid plexus cysts. *Lancet*, **346**: 724-729.
- Gupta JK, Khan KS, Thornton JG, *et al.* 1997. Management of fetal choroid plexus cysts. *Br J Obstet Gynaecol*, **104**: 881-886.
- Haddow JE, Kloza EM, Smith DE, *et al.* 1983. Data from an alpha-fetoprotein pilot screening program in Maine. *Obstet Gynecol*, **62**: 556-560.
- Haddow JE, Palomaki GE, Knight GJ, *et al.* 1998. Screening of maternal serum for fetal Down's syndrome in the first trimester. *N Engl J Med*, **338**: 955-961.
- Haddow JE, Palomaki GE, Knight GJ, Foster DL, Neveux LM. 1998. Second trimester screening for Down's syndrome using maternal serum dimeric inhibin A. *J Med Screen*, **5**: 115-119.
- Havutcu AE, Nikolopoulos G, Adinkra P, *et al.* 2002. The association between fetal pyelectasis on second trimester ultrasound scan and aneuploidy among 25,586 low risk unselected women. *Prenat Diagn*, **22**: 1201-1206.
- Hecht CA, Hook EB. 1996. Rates of Down syndrome at livebirth by one-year maternal age intervals in studies with apparent close to complete ascertainment in populations of European origin: a proposed revised rate schedule for use in genetic and prenatal screening. *Am J Med Genet*, **62**: 376-385.
- Hecht F, Bryant JS, Motulsky AG, *et al.* 1963. The No. 17-18 (E) Trisomy Syndrome. Studies on Cytogenetics, Dermatoglyphics, Paternal Age, and Linkage. *J Pediatr*, **63**: 605-621.
- Herrou M, Leporrier N, Leymarie P. 1992. Screening for fetal Down syndrome with maternal serum hCG and oestriol: a prospective study. *Prenat Diagn*, **12**: 887-892.
- Hershey DW, Crandall BF, Schroth PS. 1985. Maternal serum alpha-fetoprotein screening of fetal trisomies. *Am J Obstet Gynecol*, **153**: 224-225.
- Hertzberg BS, Kay HH, Bowie JD. 1989. Fetal choroid plexus lesions. Relationship of antenatal sonographic appearance to clinical outcome. *J Ultrasound Med*, **8**: 77-82.
- Hewitt B. 1993. Nuchal translucency in the first trimester. *Aust N Z J Obstet Gynaecol*, **33**: 389-391.
- Heyl PS, Miller W, Canick JA. 1990. Maternal serum screening for aneuploid pregnancy by alpha-fetoprotein, hCG, and unconjugated estriol. *Obstet Gynecol*, **76**: 1025-1031.
- Hogge WA, Fraer L, Melegari T. 2001. Maternal serum screening for fetal trisomy 18: benefits of patient-specific risk protocol. *Am J Obstet Gynecol*, **185**: 289-293.
- Hook EB. 1978. Spontaneous deaths of fetuses with chromosomal abnormalities diagnosed prenatally. *N Engl J Med*, **299**: 1036-1038.
- Hook EB, Topol BB, Cross PK. 1989. The natural history of cytogenetically abnormal fetuses detected at midtrimester amniocentesis which are not terminated electively: new data

- and estimates of the excess and relative risk of late fetal death associated with 47,+21 and some other abnormal karyotypes. *Am J Hum Genet*, **45**: 855-861.
- Howard RJ, Tuck SM, Long J, *et al.* 1992. The significance of choroid plexus cysts in fetuses at 18-20 weeks. An indication for amniocentesis? *Prenat Diagn*, **12**: 685-688.
- Hsu JJ, Chiang CH, Hsieh CC, *et al.* 2003. Elevated second-trimester maternal serum inhibin A levels in Asian pregnancies with fetal down syndrome and other chromosomal abnormalities. *Fetal Diagn Ther*, **18**: 105-110.
- Huderer-Duric K, Skrablin S, Kuvacic I, *et al.* 2000. The triple-marker test in predicting fetal aneuploidy: a compromise between sensitivity and specificity. *Eur J Obstet Gynecol Reprod Biol*, **88**: 49-55.
- Huether CA, Martin RL, Stoppelman SM, *et al.* 1996. Sex ratios in fetuses and liveborn infants with autosomal aneuploidy. *Am J Med Genet*, **63**: 492-500.
- Huisman TW, Bilardo CM. 1997. Transient increase in nuchal translucency thickness and reversed end-diastolic ductus venosus flow in a fetus with trisomy 18. *Ultrasound Obstet Gynecol*, **10**: 397-399.
- Hulten M 1994. Obituary: Professor Jerome Jejeune. (ed.)^(eds.).
- Hyett J, Moscoso G, Nicolaides K. 1997. Abnormalities of the heart and great arteries in first trimester chromosomally abnormal fetuses. *Am J Med Genet*, **69**: 207-216.
- Hyett JA, Sebire NJ, Snijders RJ, *et al.* 1996. Intrauterine lethality of trisomy 21 fetuses with increased nuchal translucency thickness. *Ultrasound Obstet Gynecol*, **7**: 101-103.
- Jackson LG, Zachary JM, Fowler SE, *et al.* 1992. A randomized comparison of transcervical and transabdominal chorionic-villus sampling. The U.S. National Institute of Child Health and Human Development Chorionic-Villus Sampling and Amniocentesis Study Group. *N Engl J Med*, **327**: 594-598.
- Jacobs PA, Baikie AG, Court Brown WM, *et al.* 1959. The somatic chromosomes in mongolism. *Lancet*, **1**: 710.
- Jacobson CB, Barter RH. 1967. Intrauterine diagnosis and management of genetic defects. *Am J Obstet Gynecol*, **99**: 796-807.
- Jaques AM, Halliday JL, Francis I, *et al.* 2007. Follow up and evaluation of the Victorian first-trimester combined screening programme for Down syndrome and trisomy 18. *Bjog*, **114**: 812-818.
- Jauniaux E, Nicolaides KH, Nagy AM, *et al.* 1996. Total amount of circulating human chorionic gonadotrophin alpha and beta subunits in first trimester trisomies 21 and 18. *J Endocrinol*, **148**: 27-31.
- Jelliffe-Pawlowski LL, Walton-Haynes L, Currier RJ. 2008. Using second trimester ultrasound and maternal serum biomarker data to help detect congenital heart defects in

- pregnancies with positive triple-marker screening results. *Am J Med Genet A*, **146A**: 2455-2467.
- Johnson A, Cowchock FS, Darby M, *et al.* 1991. First-trimester maternal serum alpha-fetoprotein and chorionic gonadotropin in aneuploid pregnancies. *Prenat Diagn*, **11**: 443-450.
- Johnson JM, Wilson RD, Singer J, *et al.* 1999. Technical factors in early amniocentesis predict adverse outcome. Results of the Canadian Early (EA) versus Mid-trimester (MA) Amniocentesis Trial. *Prenat Diagn*, **19**: 732-738.
- Jou HJ, Shyu MK, Lee CN, Shih JC, Guo YS, Huang CL, Hsieh FJ. 1997. Fetuses with choroid plexus cysts are at high risk for Down syndrome. *J Matern Fetal Invest*, **7**: 12-15.
- Kagan KO, Cicero S, Staboulidou I, *et al.* 2009a. Fetal nasal bone in screening for trisomies 21, 18 and 13 and Turner syndrome at 11-13 weeks of gestation. *Ultrasound Obstet Gynecol*, **33**: 259-264.
- Kagan KO, Valencia C, Livanos P, *et al.* 2009b. Tricuspid regurgitation in screening for trisomies 21, 18 and 13 and Turner syndrome at 11+0 to 13+6 weeks of gestation. *Ultrasound Obstet Gynecol*, **33**: 18-22.
- Kagan KO, Wright D, Maiz N, *et al.* 2008a. Screening for trisomy 18 by maternal age, fetal nuchal translucency, free beta-human chorionic gonadotropin and pregnancy-associated plasma protein-A. *Ultrasound Obstet Gynecol*, **32**: 488-492.
- Kagan KO, Wright D, Valencia C, *et al.* 2008b. Screening for trisomies 21, 18 and 13 by maternal age, fetal nuchal translucency, fetal heart rate, free beta-hCG and pregnancy-associated plasma protein-A. *Hum Reprod*, **23**: 1968-1975.
- Kaneko Y, Kobayashi J, Yamamoto Y, *et al.* 2008. Intensive cardiac management in patients with trisomy 13 or trisomy 18. *Am J Med Genet A*, **146A**: 1372-1380.
- Kellner LH, Weiss RR, Weiner Z, *et al.* 1995. The advantages of using triple-marker screening for chromosomal abnormalities. *Am J Obstet Gynecol*, **172**: 831-836.
- Kelly M, Robinson BW, Moore JW. 2002. Trisomy 18 in a 20-year-old woman. *Am J Med Genet*, **112**: 397-399.
- Kennedy DM, Edwards VM, Worthington DJ. 2000. Maternal serum screening for trisomy 18: assessing different statistical models to optimize detection rates. *Prenat Diagn*, **20**: 676-679.
- Kim MH, Park SH, Cho HJ, *et al.* 2006. Threshold of nuchal translucency for the detection of chromosomal aberration: comparison of different cut-offs. *J Korean Med Sci*, **21**: 11-14.
- Kim SK, Bai SW, Chung JE, *et al.* 2001. Triple marker screening for fetal chromosomal abnormalities in Korean women of advanced maternal age. *Yonsei Med J*, **42**: 199-203.

- Kiserud T. 1997. In a different vein: the ductus venosus could yield much valuable information. *Ultrasound Obstet Gynecol*, **9**: 369-372.
- Klinger K, Landes G, Shook D, *et al.* 1992. Rapid detection of chromosome aneuploidies in uncultured amniocytes by using fluorescence in situ hybridization (FISH). *Am J Hum Genet*, **51**: 55-65.
- Knight GJ, Palomaki GE, Neveux LM, *et al.* 1998. hCG and the free beta-subunit as screening tests for Down syndrome. *Prenat Diagn*, **18**: 235-245.
- Knight GJ, Palomaki GE, Neveux LM, *et al.* 2005. Integrated serum screening for Down syndrome in primary obstetric practice. *Prenat Diagn*, **25**: 1162-1167.
- Kornman LH, Morssink LP, Beekhuis JR, *et al.* 1996. Nuchal translucency cannot be used as a screening test for chromosomal abnormalities in the first trimester of pregnancy in a routine ultrasound practice. *Prenat Diagn*, **16**: 797-805.
- Kosho T. 2008. Care of children with trisomy 18 in Japan. *Am J Med Genet A*, **146A**: 1369-1371.
- Koster MP, Stoutenbeek P, Visser GH, *et al.* 2010. Trisomy 18 and 13 screening: consequences for the Dutch Down syndrome screening programme. *Prenat Diagn*, **30**: 287-289.
- Koster MP, Wortelboer EJ, Cuckle HS, *et al.* 2009. Placental protein 13 as a first trimester screening marker for aneuploidy. *Prenat Diagn*, **29**: 1237-1241.
- Kozlowski P, Knippel AJ, Froehlich S, *et al.* 2006. Additional performance of nasal bone in first trimester screening. *Ultraschall Med*, **27**: 336-339.
- Krantz DA, Hallahan TW, Orlandi F, *et al.* 2000. First-trimester Down syndrome screening using dried blood biochemistry and nuchal translucency. *Obstet Gynecol*, **96**: 207-213.
- Kratzer PG, Golbus MS, Monroe SE, *et al.* 1991. First-trimester aneuploidy screening using serum human chorionic gonadotropin (hCG), free ahCG, and progesterone. *Prenat Diagn*, **11**: 751-763.
- Kratzer PG, Golbus MS, Schonberg SA, *et al.* 1992. Cytogenetic evidence for enhanced selective miscarriage of trisomy 21 pregnancies with advancing maternal age. *Am J Med Genet*, **44**: 657-663.
- Kuhn P, Brizot ML, Pandya PP, *et al.* 1995. Crown-rump length in chromosomally abnormal fetuses at 10 to 13 weeks' gestation. *Am J Obstet Gynecol*, **172**: 32-35.
- Kupferminc MJ, Tamura RK, Sabbagha RE, *et al.* 1994. Isolated choroid plexus cyst(s): an indication for amniocentesis. *Am J Obstet Gynecol*, **171**: 1068-1071.
- Lambert-Messerlian G, Canick J, Malone F, Porter F, Nyberg D, Comstock C, Saade G, Berkowitz R, Gross S, Dugoff L, Craigo S, Timor I, Carr S, Wolfe H, D'Alton M. 2004a. First trimester levels of maternal serum hCG and inhibin A in pregnancies affected by

- fetal trisomy 18: A FASTER Trial Study. Annual Meeting of The American College of Medical Genetics, Orlando, FL, March 4-7.
- Lambert-Messerlian G, Palomaki G, Knight G, Neveux L, Canick J, Haddow J. 2004b. Dimeric inhibin-A as a marker for Down syndrome in the first trimester (Abstract #63). *Gent Med*, **6**: 277.
- Lambert-Messerlian GM, Saller DN, Jr., Tumber MB, *et al.* 1998. Second-trimester maternal serum inhibin A levels in fetal trisomy 18 and Turner syndrome with and without hydrops. *Prenat Diagn*, **18**: 1061-1067.
- Legler TJ, Lynen R, Maas JH, *et al.* 2002. Prediction of fetal Rh D and Rh CcEe phenotype from maternal plasma with real-time polymerase chain reaction. *Transfus Apher Sci*, **27**: 217-223.
- Lejeune J, Gautier M, Turpin R. 1959. [Study of somatic chromosomes from 9 mongoloid children.]. *C R Hebd Seances Acad Sci*, **248**: 1721-1722.
- Leporrier N, Herrou M, Herlicoviez M, *et al.* 1996. The usefulness of hCG and unconjugated oestriol in prenatal diagnosis of trisomy 18. *Br J Obstet Gynaecol*, **103**: 335-338.
- Leung TY, Chan LW, Leung TN, *et al.* 2007. First-trimester combined screening for trisomy 21 in a predominantly Chinese population. *Ultrasound Obstet Gynecol*, **29**: 14-17.
- Leuthner SR. 2004. Fetal palliative care. *Clin Perinatol*, **31**: 649-665.
- Levy DW, Mintz MC. 1988. The left ventricular echogenic focus: a normal finding. *AJR Am J Roentgenol*, **150**: 85-86.
- Lewis AJ. 1964. The Pathology of 18 Trisomy. *J Pediatr*, **65**: 92-101.
- Liao AW, Snijders R, Geerts L, *et al.* 2000. Fetal heart rate in chromosomally abnormal fetuses. *Ultrasound Obstet Gynecol*, **16**: 610-613.
- Lin HY, Lin SP, Chen YJ, *et al.* 2006. Clinical characteristics and survival of trisomy 18 in a medical center in Taipei, 1988-2004. *Am J Med Genet A*, **140**: 945-951.
- Lindenbaum RH, Ryyanen M, Holmes-Siedle M, *et al.* 1987. Trisomy 18 and maternal serum and amniotic fluid alpha-fetoprotein. *Prenat Diagn*, **7**: 511-519.
- Lo YM. 2001. Fetal DNA in maternal plasma: application to non-invasive blood group genotyping of the fetus. *Transfus Clin Biol*, **8**: 306-310.
- Lo YM, Corbetta N, Chamberlain PF, *et al.* 1997. Presence of fetal DNA in maternal plasma and serum. *Lancet*, **350**: 485-487.
- Lo YM, Tsui NB, Chiu RW, *et al.* 2007. Plasma placental RNA allelic ratio permits noninvasive prenatal chromosomal aneuploidy detection. *Nat Med*, **13**: 218-223.
- Lubusky M, Dhaifalah I, Prochazka M, *et al.* 2007. Single umbilical artery and its siding in the second trimester of pregnancy: relation to chromosomal defects. *Prenatal Diagnosis*, **27**: 327-331.

- Lynch L, Berkowitz RL. 1989. First trimester growth delay in trisomy 18. *Am J Perinatol*, **6**: 237-239.
- MacIntosh MC, Brambati B, Chard T, *et al.* 1993. First-trimester maternal serum Schwangerschafts protein 1 (SP1) in pregnancies associated with chromosomal anomalies. *Prenat Diagn*, **13**: 563-568.
- MacLeod AM, McHugo JM. 1991. Prenatal diagnosis of nuchal cystic hygroma. *Br J Radiol*, **64**: 802-807.
- Macri JN, Buchanan PD, Gold MP. 1986. Low alpha-fetoprotein and trisomy. *Lancet*, **2**: 405.
- Maeda T, Ohno M, Matsunobu A, *et al.* 1991. A cytogenetic survey of 14,835 consecutive liveborns. *Jinrui Idengaku Zasshi*, **36**: 117-129.
- Maiz N, Valencia C, Kagan KO, *et al.* 2009. Ductus venosus Doppler in screening for trisomies 21, 18 and 13 and Turner syndrome at 11-13 weeks of gestation. *Ultrasound Obstet Gynecol*, **33**: 512-517.
- Malone FD, Ball RH, Nyberg DA, *et al.* 2004. First-trimester nasal bone evaluation for aneuploidy in the general population. *Obstet Gynecol*, **104**: 1222-1228.
- Malone FD, Canick JA, Ball RH, *et al.* 2005. First-trimester or second-trimester screening, or both, for Down's syndrome. *N Engl J Med*, **353**: 2001-2011.
- Mansfield ES. 1993. Diagnosis of Down syndrome and other aneuploidies using quantitative polymerase chain reaction and small tandem repeat polymorphisms. *Hum Mol Genet*, **2**: 43-50.
- Martinez JM, Echevarria M, Borrell A, *et al.* 1998. Fetal heart rate and nuchal translucency in detecting chromosomal abnormalities other than Down syndrome. *Obstet Gynecol*, **92**: 68-71.
- Mastroiacovo P, Botto LD, Cavalcanti DP, *et al.* 1992. Limb anomalies following chorionic villus sampling: a registry based case-control study. *Am J Med Genet*, **44**: 856-864.
- Matias A, Gomes C, Flack N, *et al.* 1998. Screening for chromosomal abnormalities at 10-14 weeks: the role of ductus venosus blood flow. *Ultrasound Obstet Gynecol*, **12**: 380-384.
- Mavrides E, Holden D, Bland JM, *et al.* 2001. Intraobserver and interobserver variability of transabdominal Doppler velocimetry measurements of the fetal ductus venosus between 10 and 14 weeks of gestation. *Ultrasound Obstet Gynecol*, **17**: 306-310.
- Mavrides E, Sairam S, Hollis B, *et al.* 2002. Screening for aneuploidy in the first trimester by assessment of blood flow in the ductus venosus. *Bjog*, **109**: 1015-1019.
- Mazza V, Falcinelli C, Percesepe A, *et al.* 2002. Non-invasive first trimester fetal gender assignment in pregnancies at risk for X-linked recessive diseases. *Prenat Diagn*, **22**: 919-924.

- McDuffie RS, Jr., Haverkamp AD, Stark CF, *et al.* 1996. Prenatal screening using maternal serum alpha-fetoprotein, human chorionic gonadotropin, and unconjugated estriol: two-year experience in a health maintenance organization. *J Matern Fetal Med*, **5**: 70-73.
- McHugo J, Constantine G, Rajah R. 1991. Choroid plexus cysts and chromosomal defects. *Br J Radiol*, **64**: 281.
- Merkatz IR, Nitowsky HM, Macri JN, *et al.* 1984. An association between low maternal serum alpha-fetoprotein and fetal chromosomal abnormalities. *Am J Obstet Gynecol*, **148**: 886-894.
- Miell JP, Langford KS, Jones JS, *et al.* 1997. The maternal insulin-like growth factor (IGF) and IGF-binding protein response to trisomic pregnancy during the first trimester: a possible diagnostic tool for trisomy 18 pregnancies. *J Clin Endocrinol Metab*, **82**: 287-292.
- Montenegro N, Matias A, Areias JC, *et al.* 1997. Increased fetal nuchal translucency: possible involvement of early cardiac failure. *Ultrasound Obstet Gynecol*, **10**: 265-268.
- Moran CJ, Tay JB, Morrison JJ. 2002. Ultrasound detection and perinatal outcome of fetal trisomies 21, 18 and 13 in the absence of a routine fetal anomaly scan or biochemical screening. *Ultrasound Obstet Gynecol*, **20**: 482-485.
- Morcos CL, Platt LD, Carlson DE, *et al.* 1998. The isolated choroid plexus cyst. *Obstet Gynecol*, **92**: 232-236.
- Morris JK, Mutton DE, Alberman E. 2002. Revised estimates of the maternal age specific live birth prevalence of Down's syndrome. *J Med Screen*, **9**: 2-6.
- Morris JK, Savva GM. 2008. The risk of fetal loss following a prenatal diagnosis of trisomy 13 or trisomy 18. *Am J Med Genet A*, **146**: 827-832.
- MRC. 1991. Medical Research Council European trial of chorion villus sampling. MRC working party on the evaluation of chorion villus sampling. *Lancet*, **337**: 1491-1499.
- Muller F, Sault C, Lemay C, *et al.* 2002. Second trimester two-step trisomy 18 screening using maternal serum markers. *Prenat Diagn*, **22**: 605-608.
- Munson D, Leuthner SR. 2007. Palliative care for the family carrying a fetus with a life-limiting diagnosis. *Pediatr Clin North Am*, **54**: 787-798, xii.
- Murta CG, Moron AF, Avila MA, *et al.* 2002. Application of ductus venosus Doppler velocimetry for the detection of fetal aneuploidy in the first trimester of pregnancy. *Fetal Diagn Ther*, **17**: 308-314.
- Nadel A, Bromley B, Benacerraf BR. 1993. Nuchal thickening or cystic hygromas in first- and early second-trimester fetuses: prognosis and outcome. *Obstet Gynecol*, **82**: 43-48.
- Nadel AS, Bromley BS, Frigoletto FD, Jr., *et al.* 1992. Isolated choroid plexus cysts in the second-trimester fetus: is amniocentesis really indicated? *Radiology*, **185**: 545-548.

- Naguib KK, Al-Awadi SA, Moussa MA, *et al.* 1999. Trisomy 18 in Kuwait. *Int J Epidemiol*, **28**: 711-716.
- Nava S, Godmilow L, Reeser S, *et al.* 1994. Significance of sonographically detected second-trimester choroid plexus cysts: a series of 211 cases and a review of the literature. *Ultrasound Obstet Gynecol*, **4**: 448-451.
- Nawroz H, Koch W, Anker P, *et al.* 1996. Microsatellite alterations in serum DNA of head and neck cancer patients. *Nat Med*, **2**: 1035-1037.
- Nebiolo L, Ozturk M, Brambati B, *et al.* 1990. First-trimester maternal serum alpha-fetoprotein and human chorionic gonadotropin screening for chromosome defects. *Prenat Diagn*, **10**: 575-581.
- Nembhard WN, Waller DK, Sever LE, *et al.* 2001. Patterns of first-year survival among infants with selected congenital anomalies in Texas, 1995-1997. *Teratology*, **64**: 267-275.
- Ng EK, Tsui NB, Lam NY, *et al.* 2002. Presence of filterable and nonfilterable mRNA in the plasma of cancer patients and healthy individuals. *Clin Chem*, **48**: 1212-1217.
- Nicolaides KH, Azar G, Byrne D, *et al.* 1992a. Fetal nuchal translucency: ultrasound screening for chromosomal defects in first trimester of pregnancy. *Bmj*, **304**: 867-869.
- Nicolaides KH, Brizot ML, Snijders RJ. 1994. Fetal nuchal translucency: ultrasound screening for fetal trisomy in the first trimester of pregnancy. *Br J Obstet Gynaecol*, **101**: 782-786.
- Nicolaides KH, Campbell S, Gabbe SG, *et al.* 1986. Ultrasound screening for spina bifida: cranial and cerebellar signs. *Lancet*, **2**: 72-74.
- Nicolaides KH, Snijders RJ, Cuckle HS. 1998. Correct estimation of parameters for ultrasound nuchal translucency screening. *Prenat Diagn*, **18**: 519-523.
- Nicolaides KH, Snijders RJ, Gosden CM, *et al.* 1992b. Ultrasonographically detectable markers of fetal chromosomal abnormalities. *Lancet*, **340**: 704-707.
- Nicolaides KH, Spencer K, Avgidou K, *et al.* 2005. Multicenter study of first-trimester screening for trisomy 21 in 75 821 pregnancies: results and estimation of the potential impact of individual risk-orientated two-stage first-trimester screening. *Ultrasound Obstet Gynecol*, **25**: 221-226.
- Niedrist D, Riegel M, Achermann J, *et al.* 2006. Survival with trisomy 18--data from Switzerland. *Am J Med Genet A*, **140**: 952-959.
- Nielsen J, Wohlert M. 1991. Chromosome abnormalities found among 34,910 newborn children: results from a 13-year incidence study in Arhus, Denmark. *Hum Genet*, **87**: 81-83.
- Niemimaa M, Suonpaa M, Perheentupa A, *et al.* 2001. Evaluation of first trimester maternal serum and ultrasound screening for Down's syndrome in Eastern and Northern Finland. *Eur J Hum Genet*, **9**: 404-408.

- Norgaard-Pedersen B, Larsen SO, Arends J, *et al.* 1990. Maternal serum markers in screening for Down syndrome. *Clin Genet*, **37**: 35-43.
- Nyberg DA, Kramer D, Resta RG, *et al.* 1993. Prenatal sonographic findings of trisomy 18: review of 47 cases. *J Ultrasound Med*, **12**: 103-113.
- O'Callaghan SP, Giles WB, Raymond SP, *et al.* 2000. First trimester ultrasound with nuchal translucency measurement for Down syndrome risk estimation using software developed by the Fetal Medicine Foundation, United Kingdom--the first 2000 examinations in Newcastle, New South Wales, Australia. *Aust N Z J Obstet Gynaecol*, **40**: 292-295.
- Oettinger M, Odeh M, Korenblum R, *et al.* 1993. Antenatal diagnosis of choroid plexus cyst: suggested management. *Obstet Gynecol Surv*, **48**: 635-639.
- Orlandi F, Bilardo CM, Campogrande M, *et al.* 2003. Measurement of nasal bone length at 11-14 weeks of pregnancy and its potential role in Down syndrome risk assessment. *Ultrasound Obstet Gynecol*, **22**: 36-39.
- Orlandi F, Damiani G, Hallahan TW, *et al.* 1997. First-trimester screening for fetal aneuploidy: biochemistry and nuchal translucency. *Ultrasound Obstet Gynecol*, **10**: 381-386.
- Ostlere SJ, Irving HC, Lilford RJ. 1990. Fetal choroid plexus cysts: a report of 100 cases. *Radiology*, **175**: 753-755.
- Otano L, Aiello H, Igarzabal L, *et al.* 2002. Association between first trimester absence of fetal nasal bone on ultrasound and Down syndrome. *Prenat Diagn*, **22**: 930-932.
- Ozturk M, Milunsky A, Brambati B, *et al.* 1990. Abnormal maternal serum levels of human chorionic gonadotropin free subunits in trisomy 18. *Am J Med Genet*, **36**: 480-483.
- Palomaki GE, Haddow JE. 1987. Maternal serum alpha-fetoprotein, age, and Down syndrome risk. *Am J Obstet Gynecol*, **156**: 460-463.
- Palomaki GE, Haddow JE, Knight GJ, *et al.* 1995. Risk-based prenatal screening for trisomy 18 using alpha-fetoprotein, unconjugated oestriol and human chorionic gonadotropin. *Prenat Diagn*, **15**: 713-723.
- Palomaki GE, Knight G, Neveux L, Pandian R, Haddow J 2004. Maternal serum invasive trophoblast antigen (ITA) in first trimester trisomy 18 pregnancies. 2004 ASHG Annual Meeting. (ed.)^(eds.).
- Palomaki GE, Knight GJ, Haddow JE, *et al.* 1992. Prospective intervention trial of a screening protocol to identify fetal trisomy 18 using maternal serum alpha-fetoprotein, unconjugated oestriol, and human chorionic gonadotropin. *Prenat Diagn*, **12**: 925-930.
- Palomaki GE, Knight GJ, Neveux LM, *et al.* 2005. Maternal serum invasive trophoblast antigen and first-trimester Down syndrome screening. *Clin Chem*, **51**: 1499-1504.
- Palomaki GE, Lambert-Messerlian GM, Canick JA. 2007. A summary analysis of Down syndrome markers in the late first trimester. *Adv Clin Chem*, **43**: 177-210.

- Palomaki GE, Neveux LM, Donnfeld A, *et al.* 2008. Quality assessment of routine nuchal translucency measurements: a North American laboratory perspective. *Genet Med*, **10**: 131-138.
- Palomaki GE, Neveux LM, Knight GJ, *et al.* 2003. Maternal serum-integrated screening for trisomy 18 using both first- and second-trimester markers. *Prenat Diagn*, **23**: 243-247.
- Palomaki GE, Steinort K, Knight GJ, *et al.* 2006. Comparing three screening strategies for combining first- and second-trimester Down syndrome markers. *Obstet Gynecol*, **107**: 367-375.
- Palomaki GE, Williams J, Haddow JE. 1989. Combining maternal serum alpha-fetoprotein measurements and age to screen for Down syndrome in pregnant women under age 35. *Am J Obstet Gynecol*, **160**: 575-581.
- Pandya PP, Brizot ML, Kuhn P, *et al.* 1994. First-trimester fetal nuchal translucency thickness and risk for trisomies. *Obstet Gynecol*, **84**: 420-423.
- Pandya PP, Kondylis A, Hilbert L, *et al.* 1995. Chromosomal defects and outcome in 1015 fetuses with increased nuchal translucency. *Ultrasound Obstet Gynecol*, **5**: 15-19.
- Papp C, Szigeti Z, Joo JG, *et al.* 2007. The role of perinatal autopsy in the management of pregnancies with major fetal trisomies. *Pathol Res Pract*, **203**: 525-531.
- Papp C, Szigeti Z, Toth-Pal E, *et al.* 2008. Ultrasonographic findings of fetal aneuploidies in the second trimester--our experiences. *Fetal Diagn Ther*, **23**: 105-113.
- Parker MJ, Budd JL, Draper ES, *et al.* 2003. Trisomy 13 and trisomy 18 in a defined population: epidemiological, genetic and prenatal observations. *Prenat Diagn*, **23**: 856-860.
- Patau K, Smith DW, Therman E, *et al.* 1960. Multiple congenital anomaly caused by an extra autosome. *Lancet*, **1**: 790-793.
- Peralta CFA, Falcon O, Wegrzyn P, *et al.* 2005. Assessment of the gap between the fetal nasal bones at 11 to 13 + 6 weeks of gestation by three-dimensional ultrasound. *Ultrasound in Obstetrics and Gynecology*, **25**: 464-467.
- Perni SC, Predanic M, Kalish RB, *et al.* 2006. Clinical use of first-trimester aneuploidy screening in a United States population can replicate data from clinical trials. *Am J Obstet Gynecol*, **194**: 127-130.
- Perpignano MC, Cohen HL, Klein VR, *et al.* 1992. Fetal choroid plexus cysts: beware the smaller cyst. *Radiology*, **182**: 715-717.
- Petek E, Pertl B, Tschernigg M, *et al.* 2003. Characterisation of a 19-year-old "long-term survivor" with Edwards syndrome. *Genet Couns*, **14**: 239-244.
- Petrone J 2010. Q&A: Columbia's Ronald Wapner on Applying Arrah CGH in Prenatal Medicine. (ed.)^(eds.).

- Picklesimer AH, Moise KJ, Jr., Wolfe HM. 2005. The impact of gestational age on the sonographic detection of aneuploidy. *Am J Obstet Gynecol*, **193**: 1243-1247.
- Platt LD, Carlson DE, Medearis AL, *et al.* 1991. Fetal choroid plexus cysts in the second trimester of pregnancy: a cause for concern. *Am J Obstet Gynecol*, **164**: 1652-1655; discussion 1655-1656.
- Podobnik M, Singer Z, Podobnik-Sarkanji S, *et al.* 1995. First trimester diagnosis of cystic hygromata using transvaginal ultrasound and cytogenetic evaluation. *J Perinat Med*, **23**: 283-291.
- Poon LC, Chelemen T, Minekawa R, *et al.* 2009. Maternal serum ADAM12 (A disintegrin and metalloprotease) in chromosomally abnormal pregnancy at 11-13 weeks. *Am J Obstet Gynecol*, **200**: 508 e501-506.
- Porto M, Murata Y, Warneke LA, *et al.* 1993. Fetal choroid plexus cysts: an independent risk factor for chromosomal anomalies. *J Clin Ultrasound*, **21**: 103-108.
- Prefumo F, De Biasio P, Venturini PL. 2001. Reproducibility of ductus venosus Doppler flow measurements at 11-14 weeks of gestation. *Ultrasound Obstet Gynecol*, **17**: 301-305.
- Prefumo F, Sethna F, Sairam S, *et al.* 2005. First-trimester ductus venosus, nasal bones, and Down syndrome in a high-risk population. *Obstet Gynecol*, **105**: 1348-1354.
- Ramirez-Castro JL, Bersu ET. 1978. Anatomical analysis of the developmental effects of aneuploidy in man--the 18-trisomy syndrome: II. Anomalies of the upper and lower limbs. *Am J Med Genet*, **2**: 285-306.
- Rasmussen SA, Wong LY, Yang Q, *et al.* 2003. Population-based analyses of mortality in trisomy 13 and trisomy 18. *Pediatrics*, **111**: 777-784.
- Reinsch RC. 1997. Choroid plexus cysts--association with trisomy: prospective review of 16,059 patients. *Am J Obstet Gynecol*, **176**: 1381-1383.
- Rhoads GG, Jackson LG, Schlesselman SE, *et al.* 1989. The safety and efficacy of chorionic villus sampling for early prenatal diagnosis of cytogenetic abnormalities. *N Engl J Med*, **320**: 609-617.
- Ricketts NE, Lowe EM, Patel NB. 1987. Prenatal diagnosis of choroid plexus cysts. *Lancet*, **1**: 213-214.
- Ristic O, Fryns JP, Kleczowska A, *et al.* 1991. On the recurrence risk of 18 trisomy. *Ann Genet*, **34**: 47-48.
- Roberts AB, Hampton E, Wilson N. 1993. Ultrasound detection of fetal structural abnormalities in Auckland 1988-9. *N Z Med J*, **106**: 441-443.
- Root S, Carey JC. 1994. Survival in trisomy 18. *Am J Med Genet*, **49**: 170-174.
- Rosen T, D'Alton ME, Platt LD, *et al.* 2007. First-trimester ultrasound assessment of the nasal bone to screen for aneuploidy. *Obstet Gynecol*, **110**: 399-404.

- Ryynanen M, Leskinen S, Heinonen S, *et al.* 1997. Recurrence risk of a serious, noninherited chromosomal abnormality. *Fertil Steril*, **68**: 439-442.
- Saller DN, Jr., Keene CL, Sun CC, *et al.* 1990. The association of single umbilical artery with cytogenetically abnormal pregnancies. *Am J Obstet Gynecol*, **163**: 922-925.
- Sancken U, Bartels I, Louwen F, *et al.* 1999. A retrospective evaluation of second-trimester serum screening for fetal trisomy 18: experience of two laboratories. *Prenat Diagn*, **19**: 947-954.
- Savoldelli G, Binkert F, Achermann J, *et al.* 1993. Ultrasound screening for chromosomal anomalies in the first trimester of pregnancy. *Prenat Diagn*, **13**: 513-518.
- Savva GM, Walker K, Morris JK. 2010. The maternal age-specific live birth prevalence of trisomies 13 and 18 compared to trisomy 21 (Down syndrome). *Prenat Diagn*, **30**: 57-64.
- Schaelike M, Kossakiewicz M, Kossakiewicz A, *et al.* 2009. Examination of a first-trimester Down syndrome screening concept on a mix of 11,107 high- and low-risk patients at a private center for prenatal medicine in Germany. *Eur J Obstet Gynecol Reprod Biol*, **144**: 140-145.
- Schneider AS, Mennuti MT, Zackai EH. 1981. High cesarean section rate in trisomy 18 births: a potential indication for late prenatal diagnosis. *Am J Obstet Gynecol*, **140**: 367-370.
- Schuchter K, Hafner E, Stangl G, *et al.* 2001. Sequential screening for trisomy 21 by nuchal translucency measurement in the first trimester and maternal serum biochemistry in the second trimester in a low-risk population. *Ultrasound Obstet Gynecol*, **18**: 23-25.
- Scott F, Wheeler D, Sinosich M, *et al.* 1996. First trimester aneuploidy screening using nuchal translucency, free beta human chorionic gonadotrophin and maternal age. *Aust N Z J Obstet Gynaecol*, **36**: 381-384.
- Senat MV, Bernard JP, Boulvain M, *et al.* 2003. Intra- and interoperator variability in fetal nasal bone assessment at 11-14 weeks of gestation. *Ultrasound Obstet Gynecol*, **22**: 138-141.
- Seoud MA, Alley DC, Smith DL, *et al.* 1994. Prenatal sonographic findings in trisomy 13, 18, 21 and 22. A review of 46 cases. *J Reprod Med*, **39**: 781-787.
- Seppo H, Markku R, Marko N, *et al.* 1999. Detection of trisomy 18 by double screening in a low-risk pregnant population. *Fetal Diagn Ther*, **14**: 15-19.
- Sepulveda W, Cullen S, Nicolaidis P, *et al.* 1995. Echogenic foci in the fetal heart: a marker of chromosomal abnormality. *Br J Obstet Gynaecol*, **102**: 490-492.
- Sepulveda W, Wong AE, Dezerega V. 2010. First-trimester sonographic findings in trisomy 18: a review of 53 cases. *Prenat Diagn*, **30**: 256-259.
- Shaffer LG, Bui TH. 2007. Molecular cytogenetic and rapid aneuploidy detection methods in prenatal diagnosis. *Am J Med Genet C Semin Med Genet*, **145C**: 87-98.

- Shanske AL. 2006. Trisomy 18 in a second 20-year-old woman. *Am J Med Genet A*, **140**: 966-967.
- Sherod C, Sebire NJ, Soares W, *et al.* 1997. Prenatal diagnosis of trisomy 18 at the 10-14-week ultrasound scan. *Ultrasound Obstet Gynecol*, **10**: 387-390.
- Shields LE, Carpenter LA, Smith KM, *et al.* 1998. Ultrasonographic diagnosis of trisomy 18: is it practical in the early second trimester? *J Ultrasound Med*, **17**: 327-331.
- Shulman LP, Emerson DS, Felker RE, *et al.* 1992. High frequency of cytogenetic abnormalities in fetuses with cystic hygroma diagnosed in the first trimester. *Obstet Gynecol*, **80**: 80-82.
- Shulman LP, Simpson J. 2008. *Glob. Libr. women's med.*, (ed.)^(eds.).
- Sifakis S, Akolekar R, Syngelaki A, De Cruz J, Nicolaides KH. 2010. Maternal serum human placental growth hormone at 11 to 13 weeks in trisomy 21 and trisomy 18 pregnancies. *Prenat Diagn*, **30**: 212-215.
- Simpson JL, Elias S. 1993. Isolating fetal cells from maternal blood. Advances in prenatal diagnosis through molecular technology. *Jama*, **270**: 2357-2361.
- Smidt-Jensen S, Permin M, Philip J, *et al.* 1992. Randomised comparison of amniocentesis and transabdominal and transcervical chorionic villus sampling. *Lancet*, **340**: 1237-1244.
- Smith DW. 1964. Autosomal Abnormalities. *Am J Obstet Gynecol*, **90**: SUPPL:1055-1077.
- Smith DW, Patau K, Therman E, Inhorn SL. 1960. A new autosomal trisomy syndrome: multiple congenital anomalies caused by an extra chromosome. *J Pediatr*, **57**: 338-345.
- Snijders RJ, Holzgreve W, Cuckle H, *et al.* 1994. Maternal age-specific risks for trisomies at 9-14 weeks' gestation. *Prenat Diagn*, **14**: 543-552.
- Snijders RJ, Sundberg K, Holzgreve W, *et al.* 1999. Maternal age- and gestation-specific risk for trisomy 21. *Ultrasound Obstet Gynecol*, **13**: 167-170.
- Spencer K. 2001. Age related detection and false positive rates when screening for Down's syndrome in the first trimester using fetal nuchal translucency and maternal serum free beta hCG and PAPP-A. *Bjog*, **108**: 1043-1046.
- Spencer K, Aitken DA, Crossley JA, *et al.* 1994. First trimester biochemical screening for trisomy 21: the role of free beta hCG, alpha fetoprotein and pregnancy associated plasma protein A. *Ann Clin Biochem*, **31 (Pt 5)**: 447-454.
- Spencer K, Bindra R, Nix AB, *et al.* 2003. Delta-NT or NT MoM: which is the most appropriate method for calculating accurate patient-specific risks for trisomy 21 in the first trimester? *Ultrasound Obstet Gynecol*, **22**: 142-148.
- Spencer K, Cowans NJ, Stamatopoulou A. 2007. Maternal serum ADAM12s as a marker of rare aneuploidies in the first or second trimester of pregnancy. *Prenat Diagn*, **27**: 1233-1237.

- Spencer K, Crossley JA, Green K, *et al.* 1999. Second trimester levels of pregnancy associated plasma protein-A in cases of trisomy 18. *Prenat Diagn*, **19**: 1127-1134.
- Spencer K, Heath V, Flack N, *et al.* 2000a. First trimester maternal serum AFP and total hCG in aneuploidies other than trisomy 21. *Prenat Diagn*, **20**: 635-639.
- Spencer K, Liao AW, Ong CY, *et al.* 2001a. Maternal serum activin A and inhibin A in trisomy 18 pregnancies at 10-14 weeks. *Prenat Diagn*, **21**: 571-574.
- Spencer K, Liao AW, Ong CY, *et al.* 2001b. First trimester maternal serum placenta growth factor (PIGF) concentrations in pregnancies with fetal trisomy 21 or trisomy 18. *Prenat Diagn*, **21**: 718-722.
- Spencer K, Macri JN, Aitken DA, *et al.* 1992. Free beta-hCG as first-trimester marker for fetal trisomy. *Lancet*, **339**: 1480.
- Spencer K, Mallard AS, Coombes EJ, *et al.* 1993. Prenatal screening for trisomy 18 with free beta human chorionic gonadotrophin as a marker. *Bmj*, **307**: 1455-1458.
- Spencer K, Noble P, Snijders RJ, *et al.* 1997. First-trimester urine free beta hCG, beta core, and total oestriol in pregnancies affected by Down's syndrome: implications for first-trimester screening with nuchal translucency and serum free beta hCG. *Prenat Diagn*, **17**: 525-538.
- Spencer K, Ong C, Skentou H, *et al.* 2000b. Screening for trisomy 13 by fetal nuchal translucency and maternal serum free beta-hCG and PAPP-A at 10-14 weeks of gestation. *Prenat Diagn*, **20**: 411-416.
- Spencer K, Spencer CE, Power M, *et al.* 2000c. One stop clinic for assessment of risk for fetal anomalies: a report of the first year of prospective screening for chromosomal anomalies in the first trimester. *Bjog*, **107**: 1271-1275.
- Staples AJ, Robertson EF, Ranieri E, *et al.* 1991. A maternal serum screen for trisomy 18: an extension of maternal serum screening for Down syndrome. *Am J Hum Genet*, **49**: 1025-1033.
- Stempfle N, Hutten Y, Fredouille C, *et al.* 1999. Skeletal abnormalities in fetuses with Down's syndrome: a radiographic post-mortem study. *Pediatr Radiol*, **29**: 682-688.
- Stene J, Stene E, Mikkelsen M. 1984. Risk for chromosome abnormality at amniocentesis following a child with a non-inherited chromosome aberration. A European Collaborative Study on Prenatal Diagnoses 1981. *Prenat Diagn*, **4 Spec No**: 81-95.
- Stenhouse EJ, Crossley JA, Aitken DA, *et al.* 2004. First-trimester combined ultrasound and biochemical screening for Down syndrome in routine clinical practice. *Prenat Diagn*, **24**: 774-780.
- Subramanian R, Sadasivan R, Rao MR, Reddy SB. 1988. An association between maternal serum alpha-fetoprotein and fetal autosomal trisomies. *Am J Hum Genet*, **43**: a250.

- Suchy SF, Yeager MT. 1990. Down syndrome screening in women under 35 with maternal serum hCG. *Obstet Gynecol*, **76**: 20-24.
- Summers AM, Farrell SA, Huang T, *et al.* 2003. Maternal serum screening in Ontario using the triple marker test. *J Med Screen*, **10**: 107-111.
- Suzumori K, Tanemura M, Murakami I, *et al.* 1997. A retrospective evaluation of maternal serum screening for the detection of fetal aneuploidy. *Prenat Diagn*, **17**: 861-866.
- Szabo J, Gellen J. 1990. Nuchal fluid accumulation in trisomy-21 detected by vaginosonography in first trimester. *Lancet*, **336**: 1133.
- Tabor A, Philip J, Bang J, *et al.* 1988. Needle size and risk of miscarriage after amniocentesis. *Lancet*, **1**: 183-184.
- Tabor A, Philip J, Madsen M, *et al.* 1986. Randomised controlled trial of genetic amniocentesis in 4606 low-risk women. *Lancet*, **1**: 1287-1293.
- Taipale P, Hiilesmaa V, Salonen R, *et al.* 1997. Increased nuchal translucency as a marker for fetal chromosomal defects. *N Engl J Med*, **337**: 1654-1658.
- Taslimi MM, Acosta R, Chueh J, *et al.* 2005. Detection of sonographic markers of fetal aneuploidy depends on maternal and fetal characteristics. *J Ultrasound Med*, **24**: 811-815.
- Taylor AI. 1968. Autosomal trisomy syndromes: a detailed study of 27 cases of Edwards' syndrome and 27 cases of Patau's syndrome. *J Med Genet*, **5**: 227-252.
- Taylor AI, Moores EC. 1967. A sex chromatin survey of newborn children in two London hospitals. *J Med Genet*, **4**: 258-259.
- Theodoropoulos P, Lolis D, Papageorgiou C, *et al.* 1998. Evaluation of first-trimester screening by fetal nuchal translucency and maternal age. *Prenat Diagn*, **18**: 133-137.
- Thorpe-Beeston JG, Gosden CM, Nicolaidis KH. 1990. Choroid plexus cysts and chromosomal defects. *Br J Radiol*, **63**: 783-786.
- Tong YK, Ding C, Chiu RW, *et al.* 2006. Noninvasive prenatal detection of fetal trisomy 18 by epigenetic allelic ratio analysis in maternal plasma: Theoretical and empirical considerations. *Clin Chem*, **52**: 2194-2202.
- Tongsong T, Sirichotiyakul S, Wanapirak C, *et al.* 2002. Sonographic features of trisomy 18 at midpregnancy. *J Obstet Gynaecol Res*, **28**: 245-250.
- Torrington N. 2009. Performance of first-trimester screening between gestational weeks 7 and 13. *Clin Chem*, **55**: 1564-1567.
- Toyama JM, Brizot ML, Liao AW, *et al.* 2004. Ductus venosus blood flow assessment at 11 to 14 weeks of gestation and fetal outcome. *Ultrasound Obstet Gynecol*, **23**: 341-345.
- Trauffer PM, Anderson CE, Johnson A, *et al.* 1994. The natural history of euploid pregnancies with first-trimester cystic hygromas. *Am J Obstet Gynecol*, **170**: 1279-1284.

- Tsai MS, Huang YY, Hwa KY, *et al.* 2001. Combined measurement of fetal nuchal translucency, maternal serum free beta-hCG, and pregnancy-associated plasma protein A for first-trimester Down's syndrome screening. *J Formos Med Assoc*, **100**: 319-325.
- Tucker ME, Garringer HJ, Weaver DD. 2007. Phenotypic spectrum of mosaic trisomy 18: two new patients, a literature review, and counseling issues. *Am J Med Genet A*, **143**: 505-517.
- Tul N, Spencer K, Noble P, *et al.* 1999. Screening for trisomy 18 by fetal nuchal translucency and maternal serum free beta-hCG and PAPP-A at 10-14 weeks of gestation. *Prenat Diagn*, **19**: 1035-1042.
- Twining P, Zuccollo J, Clewes J, *et al.* 1991. Fetal choroid plexus cysts: a prospective study and review of the literature. *Br J Radiol*, **64**: 98-102.
- Uehara S, Yaegashi N, Maeda T, *et al.* 1999. Risk of recurrence of fetal chromosomal aberrations: analysis of trisomy 21, trisomy 18, trisomy 13, and 45,X in 1,076 Japanese mothers. *J Obstet Gynaecol Res*, **25**: 373-379.
- Van Dyke DC, Allen M. 1990. Clinical management considerations in long-term survivors with trisomy 18. *Pediatrics*, **85**: 753-759.
- van Dyke DI. 2010. Amniotic Fluid Cell Culture. In *Genetic Disorders and the Fetus: Diagnosis, Prevention and Treatment*, Milunsky A, Milunsky JM (eds.). 6th ed. Wiley-Blackwell: West Sussex; 138-159.
- Van Lith JM. 1992. First-trimester maternal serum human chorionic gonadotrophin as a marker for fetal chromosomal disorders. The Dutch Working Party on Prenatal Diagnosis. *Prenat Diagn*, **12**: 495-504.
- van Zalen-Sprock RM, van Vugt JM, van Geijn HP. 1992. First-trimester diagnosis of cystic hygroma--course and outcome. *Am J Obstet Gynecol*, **167**: 94-98.
- Ville Y, Lalondrelle C, Doumerc S, *et al.* 1992. First-trimester diagnosis of nuchal anomalies: significance and fetal outcome. *Ultrasound Obstet Gynecol*, **2**: 314-316.
- Viora E, Masturzo B, Errante G, *et al.* 2003. Ultrasound evaluation of fetal nasal bone at 11 to 14 weeks in a consecutive series of 1906 fetuses. *Prenat Diagn*, **23**: 784-787.
- Viora E, Zamboni C, Mortara G, *et al.* 2007. Trisomy 18: Fetal ultrasound findings at different gestational ages. *Am J Med Genet A*, **143**: 553-557.
- von Kaisenberg CS, Gasiorek-Wiens A, Bielicki M, *et al.* 2002. Screening for trisomy 21 by maternal age, fetal nuchal translucency and maternal serum biochemistry at 11-14 weeks: a German multicenter study. *J Matern Fetal Neonatal Med*, **12**: 89-94.
- Wald NJ. 1976. The detection of neural-tube defects by screening maternal blood. In *Prenatal Diagnosis*, Boue A (eds.). Les Colloques d'INSERM; 227-238.
- Wald NJ. 2008. Guidance on terminology. *J Med Screen*, **15**: 50.

- Wald NJ, Cuckle H, Brock JH, *et al.* 1977. Maternal serum-alpha-fetoprotein measurement in antenatal screening for anencephaly and spina bifida in early pregnancy. Report of U.K. collaborative study on alpha-fetoprotein in relation to neural-tube defects. *Lancet*, **1**: 1323-1332.
- Wald NJ, Cuckle HS, Boreham J, *et al.* 1979. Antenatal screening in Oxford for fetal neural tube defects. *Br J Obstet Gynaecol*, **86**: 91-100.
- Wald NJ, Hackshaw AK, George LM. 2000. Assay precision of serum alpha fetoprotein in antenatal screening for neural tube defects and Down's syndrome. *J Med Screen*, **7**: 74-77.
- Wald NJ, Rodeck C, Hackshaw AK, *et al.* 2003. First and second trimester antenatal screening for Down's syndrome: the results of the Serum, Urine and Ultrasound Screening Study (SURUSS). *J Med Screen*, **10**: 56-104.
- Wald NJ, Rodeck C, Hackshaw AK, Walters J, Chitty L, Mackinson AM, Bestwick JP. 2006. Errata. *J Med Screen*, **13**: 51-52.
- Wald NJ, Watt HC, Hackshaw AK. 1999. Integrated screening for Down's syndrome on the basis of tests performed during the first and second trimesters. *N Engl J Med*, **341**: 461-467.
- Walkinshaw S, Pilling D, Spriggs A. 1994. Isolated choroid plexus cysts--the need for routine offer of karyotyping. *Prenat Diagn*, **14**: 663-667.
- Walknowska J, Conte FA, Grumbach MM. 1969. Practical and theoretical implications of fetal-maternal lymphocyte transfer. *Lancet*, **1**: 1119-1122.
- Wapner R, Thom E, Simpson JL, *et al.* 2003. First-trimester screening for trisomies 21 and 18. *N Engl J Med*, **349**: 1405-1413.
- Warburton D, Dallaire L, Thangavelu M, *et al.* 2004. Trisomy recurrence: a reconsideration based on North American data. *Am J Hum Genet*, **75**: 376-385.
- Watanabe H, Hamada H, Yamada N, *et al.* 2002. Second-trimester maternal pregnancy-associated plasma protein a and inhibin a levels in fetal trisomies. *Fetal Diagn Ther*, **17**: 137-141.
- Watson WJ, Miller RC, Menard MK, *et al.* 1994. Ultrasonographic measurement of fetal nuchal skin to screen for chromosomal abnormalities. *Am J Obstet Gynecol*, **170**: 583-586.
- Watson WJ, Miller RC, Wax JR, *et al.* 2008. Sonographic findings of trisomy 18 in the second trimester of pregnancy. *J Ultrasound Med*, **27**: 1033-1038; quiz 1039-1040.
- Wax JR, Royer D, Mather J, *et al.* 2000. A preliminary study of sonographic grading of fetal intracardiac echogenic foci: feasibility, reliability and association with aneuploidy. *Ultrasound Obstet Gynecol*, **16**: 123-127.

- Wayda K, Kereszturi A, Orvos H, *et al.* 2001. Four years experience of first-trimester nuchal translucency screening for fetal aneuploidies with increasing regional availability. *Acta Obstet Gynecol Scand*, **80**: 1104-1109.
- Weber WW. 1967. Survival and the sex ratio in trisomy 17-18. *Am J Hum Genet*, **19**: 369-377.
- Weingertner AS, Kohler M, Firtion C, *et al.* 2006. Interest of foetal nasal bone measurement at first trimester trisomy 21 screening. *Fetal Diagn Ther*, **21**: 433-438.
- Wenstrom KD, Chu DC, Owen J, *et al.* 1998. Maternal serum alpha-fetoprotein and dimeric inhibin A detect aneuploidies other than Down syndrome. *Am J Obstet Gynecol*, **179**: 966-970.
- WHO/PAHO. 1999. Evaluation of chorionic villus sampling safety: WHO/PAHO consultation on CVS. *Prenat Diagn*, **19**: 97-99.
- Wojdemann KR, Shalmi AC, Christiansen M, *et al.* 2005. Improved first-trimester Down syndrome screening performance by lowering the false-positive rate: a prospective study of 9941 low-risk women. *Ultrasound Obstet Gynecol*, **25**: 227-233.
- Won RH, Currier RJ, Lorey F, *et al.* 2005. The timing of demise in fetuses with trisomy 21 and trisomy 18. *Prenat Diagn*, **25**: 608-611.
- Wong SF, Choi H, Ho LC. 2003. Nasal bone hypoplasia: is it a common finding amongst chromosomally normal fetuses of southern Chinese women? *Gynecol Obstet Invest*, **56**: 99-101.
- Wortelboer EJ, Koster MP, Stoutenbeek P, *et al.* 2008. Fifteen years of triple tests in The Netherlands; the life cycle of a screening test. *Prenat Diagn*, **28**: 950-955.
- Wright D, Kagan KO, Molina FS, *et al.* 2008. A mixture model of nuchal translucency thickness in screening for chromosomal defects. *Ultrasound Obstet Gynecol*, **31**: 376-383.
- Yamanaka M, Setoyama T, Igarashi Y, *et al.* 2006. Pregnancy outcome of fetuses with trisomy 18 identified by prenatal sonography and chromosomal analysis in a perinatal center. *Am J Med Genet A*, **140**: 1177-1182.
- Yankowitz J, Fulton A, Williamson R, *et al.* 1998. Prospective evaluation of prenatal maternal serum screening for trisomy 18. *Am J Obstet Gynecol*, **178**: 446-450.
- Yeo L, Guzman ER, Day-Salvatore D, *et al.* 2003. Prenatal detection of fetal trisomy 18 through abnormal sonographic features. *J Ultrasound Med*, **22**: 581-590; quiz 591-582.
- Yoshida K, Kuwabara Y, Tanaka T, *et al.* 2000. Dimeric inhibin A as a fourth marker for Down's syndrome maternal serum screening in native Japanese women. *J Obstet Gynaecol Res*, **26**: 171-174.
- Young ID, Cook JP, Mehta L. 1986. Changing demography of trisomy 18. *Arch Dis Child*, **61**: 1035-1036.

- Zaragoza E, Akolekar R, Poon LC, *et al.* 2009. Maternal serum placental growth factor at 11-13 weeks in chromosomally abnormal pregnancies. *Ultrasound Obstet Gynecol*, **33**: 382-386.
- Zeitune M, Aitken DA, Crossley JA, *et al.* 1991. Estimating the risk of a fetal autosomal trisomy at mid-trimester using maternal serum alpha-fetoprotein and age: a retrospective study of 142 pregnancies. *Prenat Diagn*, **11**: 847-857.
- Zimmermann R, Hucha A, Savoldelli G, *et al.* 1996. Serum parameters and nuchal translucency in first trimester screening for fetal chromosomal abnormalities. *Br J Obstet Gynaecol*, **103**: 1009-1014.
- Zoppi MA, Ibba RM, Axiana C, *et al.* 2003. Absence of fetal nasal bone and aneuploidies at first-trimester nuchal translucency screening in unselected pregnancies. *Prenat Diagn*, **23**: 496-500.
- Zoppi MA, Putzolu M, Ibba RM, *et al.* 2002. First-trimester ductus venosus velocimetry in relation to nuchal translucency thickness and fetal karyotype. *Fetal Diagn Ther*, **17**: 52-57.